

eXtra Botany

Viewpoint

To tolerate drought or resist aphids? A new challenge to plant science is on the horizon

Claudio C. Ramírez^{1,*}, Pedro E. Gundel^{2,3}, Alison J. Karley⁴ and Daniel J. Leybourne⁵

¹ Centro de Ecología Molecular y Funcional, Instituto de Ciencias Biológicas, Universidad de Talca, Chile

² Centro de Ecología Integrativa, Instituto de Ciencias Biológicas, Universidad de Talca, Chile

³ IFEVA, CONICET, Facultad de Agronomía, Universidad de Buenos Aires, Buenos Aires, Argentina

⁴ Ecological Sciences Department, The James Hutton Institute, Dundee, UK

⁵ Department of Evolution, Ecology, and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK

* Correspondence: clramirez@utalca.cl

Received 16 September 2022; Editorial decision 4 January 2023; Accepted 9 January 2023

Editor: Christine Foyer, University of Birmingham, UK

Aphids are important herbivorous insects that can cause significant crop damage, leading to yield reduction and economic loss. One avenue being explored to reduce aphid impacts is the development of aphid-resistant plants. Under projected climate scenarios, it is expected that plants will be exposed to greater biotic and abiotic stress, including increased herbivorous insect infestation and exposure to prolonged periods of environmental stress, particularly drought. In response to these projections, plant–aphid interactions under drought conditions have been a subject of growing interest; however, few studies have looked at the impact of drought stress on plant resistance to aphids despite the potential importance for plant breeding. Here, we examine the latest scientific advances regarding variation in plant resistance to aphids under drought, emphasizing underlying mechanisms and functional trade-offs and propose a conceptual model relating plant tolerance to drought with plant resistance to aphids.

Plants are simultaneously subjected to multiple biotic and abiotic threats. Understanding how plants respond to these factors is essential for predicting the performance of crops, especially in response to climate change (Bellard *et al.*, 2012). In nature, plant populations are shaped by environmental conditions that select for resistance to specific factors. Additionally, strong selection

for resistance to one factor can be associated with susceptibility to another (i.e. trade-offs) (Hermes and Mattson, 1992). Similar outcomes occur during plant breeding (Denison, 2012), where selection for high yields can come at a cost of increased susceptibility to environmental stressors, or where selection for resistance traits compromises plant tolerance of other stressors. A better understanding of these phenomena is needed to predict the consequences of stress-driven trait selection in natural vegetation or crops by examining potential trade-offs in breeding for biotic stress (e.g. pest and disease resistance) versus those conferring climate resilience (e.g. drought tolerance).

Aphids (Hemiptera: Aphididae) are phytophagous insects with worldwide distribution, representing an important agricultural pest of many crops (Dixon, 1998). Aphids cause plant damage both directly and indirectly. Direct damage results from sap removal during aphid feeding. Indirect damage is caused by the transmission of plant viruses and reduced quality due to build-up of aphid honeydew which favours the growth of microbes such as sooty moulds. Climate projections have estimated both positive and negative effects of climate change on herbivorous species, although most scenarios predict that proliferation of herbivorous insects will increase worldwide (Schneider *et al.*, 2022). One potential consequence of climate change is increased drought. Prolonged periods of drought affect plant homeostasis and the interaction with other organisms and, consequently, the plant response to herbivorous insects

such as aphids (Luo and Gilbert, 2022). Because there is usually a trade-off between traits, plant breeding programmes may encounter difficulties in simultaneously improving drought tolerance and pest resistance.

A conceptual model recently proposed by Leybourne *et al.* (2021), and supported by experimental results in cereals (Kansman *et al.*, 2022; Leybourne *et al.*, 2022), suggests that plant resistance to aphids increases as water availability decreases. However, the model lacks explicit consideration of how plants differing in tolerance to drought (and thus in susceptibility to water availability) might also vary in resistance to aphids. Here, we advance this model by incorporating an evolutionary perspective which considers the variation among plant genotypes in intrinsic tolerance to drought, which has been investigated by only a few studies (Quandahor *et al.*, 2019).

Towards an aphid-plant resistance hypothesis

A recent meta-analysis by Leybourne *et al.* (2021) focused on aphid responses to drought and identified significant knowledge gaps in our understanding of the effect of drought stress on aphid-susceptible and aphid-resistant plants: only four studies compared the effect of reduced water availability on plants that are resistant and susceptible to aphids. These studies suggest that aphid performance is reduced by drought on both aphid-susceptible and aphid-resistant plants but with a stronger effect on the former (Leybourne *et al.*, 2021). To explain this, Leybourne *et al.* (2021) proposed the ‘plant resistance hypothesis’ (Fig. 1A), which predicts that lower water availability causes a differential change in chemical and mo-

lecular defences between susceptible and resistant plants. In other words, susceptible plants display a more distinctive change in plant defences along a water availability gradient than resistant plants, since resistant plants have higher basal levels of defences and a narrower range of responses (Leybourne *et al.*, 2021). This hypothesis focuses on the variation in the concentration of plant defences due to water availability, whereas water availability also affects other plant traits (Kansman *et al.*, 2022). More importantly, this hypothesis assumes that either aphid-susceptible or aphid-resistant plants do not vary in their level of drought tolerance. Plant genetic variation in the ability to resist or recover from drought might alter plant responses to short-term changes in water availability. Note that while water availability is an environmental condition, drought tolerance describes the ability of plants to resist and be resilient to (recover from) low water conditions (Tardieu, 2022). Surprisingly little is known about how plants with different levels of tolerance to drought also differ in their ability to resist aphids.

Plant tolerance to aphids under drought stress

In their relationship with aphids, plants may not only evolve resistance as antagonistic response mechanisms, but may also develop tolerance to aphids. This is another missing link within the proposed plant resistance hypothesis (Leybourne *et al.*, 2021) resulting from a lack of available research. Unlike resistance, tolerance is the ability of plants to recover from herbivore damage through growth and compensatory physiological processes (Strauss and Agrawal, 1999). Most of the evidence suggests that tolerance of and resistance to herbivores




Fig. 1. (A) Original model proposed by Leybourne *et al.* (2021) that relates the performance of aphids on aphid-resistant and aphid-susceptible plants as a function of the water availability. (B) Model 1 proposed herein relating the resistance to aphids as a function of plant drought tolerance. (C) Model 2 proposed herein results from subjecting and not subjecting different plant genotypes of a crop to drought. Each dot (blue or white) in (B) and (C) represents hypothetical different plant genotypes or intraspecific plant variants (e.g. accessions, cultivars, or varieties) for which aphid resistance and drought tolerance are estimated. For example, the herbivory resistance level of a given plant genotype is estimated as plant biomass in aphid-challenged plants versus plant biomass in control plants (not challenged by aphids), all of them grown under no water restriction. In contrast, the level of drought tolerance for a given plant genotype is estimated as plant biomass in water-stressed plants versus plant biomass in control plants (with no water restriction). Traits for future focus include drought tolerance traits that reduce aphid fitness (1) and aphid resistance traits that reduce water loss (2), particularly when trait expression is elevated under reduced water availability (3).

represent independent plant defence strategies (Pearse *et al.*, 2017). The evolution of cardenolides and regrowth ability in milkweeds is a good example (Agrawal and Fishbein, 2008), and resistance and tolerance tend to be positively correlated in crops (Leimu and Koricheva, 2006). However, plant tolerance as a defence mechanism has received little attention in aphid–plant interactions (Peterson *et al.*, 2017), and much less in relation to drought (Mitchell *et al.*, 2016). Further research is needed to assess whether plants can display cross-tolerance to drought and aphid attacks (Foyer *et al.*, 2016).

Examining interactions between drought tolerance and aphid resistance from a trade-off perspective

Plants often show trade-offs between different functions that can be explained by resource limitations and by developmental constraints at the molecular level that regulate those trade-offs (Herms and Mattson, 1992). Limited resource availability can lead to conflicting demands among different fitness-related traits, preventing plants from investing simultaneously in growth, reproduction, and defence. A negative correlation

between resistance to aphids and the ability to tolerate drought among a set of plant genotypes would indicate that tolerance to drought requires the allocation of resources for an improved water economy at the expense of defence against aphids (Fig. 1B, Model 1).

Plant genotypes could differ in the resistance level to aphids based on their intrinsic level of drought tolerance. This raises questions about the predicted responses of drought-tolerant and drought-susceptible plant genotypes to aphid attack when exposed to a water availability gradient (as in Leybourne *et al.*, 2021). Under drought conditions, do drought-susceptible plants show relatively larger increases in aphid resistance than drought-tolerant plants because the latter invest more in tolerating drought (Fig. 1C)? We propose that future studies dealing with drought and aphid attack should focus on drought tolerance traits that reduce aphid fitness and aphid resistance traits that reduce water loss, particularly when trait expression is elevated under reduced water availability (Fig. 1C). Alternatively, if resistance to aphids is independent of drought tolerance, aphid resistance might not vary under drought conditions. Disentangling these relationships is key to guiding plant domestication programmes in the context of developing climate-resilient crops.


Box 1. Potential mechanisms underpinning the interactions between drought tolerance and aphid resistance.

Plant traits conferring drought tolerance can also confer aphid resistance, and vice versa. Additionally, there is growing evidence for crosstalk between molecular signalling pathways responding to these two stressors that may explain their interaction. Cross-tolerance could result, therefore, from biochemical responses that influence osmotic potential and nutritional quality (A), physical characteristics that alter water loss and aphid infestation (B), and elevated molecular defences (C). These can also be involved in cross-tolerance and crosstalk with aphid resistance traits.

(A) Biochemical traits: osmoprotective mechanisms include changes in the composition and concentrations of secondary metabolites, soluble proteins, amino acids, and carbohydrates (Osakabe *et al.*, 2014). Concentrations of non-structural carbohydrates (NSCs) are modulated by drought and act as a carbohydrate reserve for stress (Sadras *et al.*, 2021); recently, NSCs were also reported to contribute towards plant resistance to aphids in cereals (Sadras *et al.*, 2020). Other osmoprotective metabolites, such as essential amino acids, have also been associated with aphid resistance (Leybourne *et al.*, 2019). The potential mechanism(s) through which these metabolites provide cross-tolerance against aphids could be through the low osmotic potential generated by high metabolite concentration, and reduced phloem nitrogen quality, which can limit aphid performance (Sadras *et al.*, 2020, 2021).

(B) Morphological traits: the morphological traits trichomes and epicuticular waxes can provide drought tolerance by limiting transpiration. Recent research has indicated that drought stress can stimulate the production of these physical traits (Saska *et al.*, 2021, 2022), which have also been associated with increased aphid resistance (Valim *et al.*, 2016).

(C) Defence signalling pathways: benzoxazinoids represent a key example of crosstalk between drought tolerance and resistance to aphids. The role of benzoxazinoids as defensive metabolites in aphid resistance in cereals has been well documented (Niemeyer, 2009), and linked to resistance mechanisms such as induction of callose deposition (Zhou *et al.*, 2018). Recent research has shown that benzoxazinoid biosynthesis is regulated by the drought-induced transcription factor MYB31 (Batyrschina *et al.*, 2022), indicating that it could also respond to drought. The regulation of thionin gene expression is another example of crosstalk since its expression was greater in aphid-resistant than in susceptible plants (Escudero-Martinez *et al.*, 2017; Leybourne *et al.*, 2019) and did not change in response to drought, whereas expression was up-regulated in susceptible plants (Leybourne *et al.*, 2022). The role of these metabolites in drought tolerance has yet to be established.



A mechanistic approach to understand the relationship between aphid resistance and drought tolerance

Plant resistance to aphids can be conferred by chemical deterrence traits, physical barriers to aphid settling and feeding, and traits that reduce plant quality for feeding (Mitchell *et al.*, 2016). Plant traits conferring tolerance to drought include the accumulation of metabolites that maintain turgor and tissue functionality under water scarcity (Benkeblia, 2022), mechanisms to regulate stomatal aperture and tissue relative water content (Buckley, 2019), and changes to root and leaf tissue structure (Fang and Xiong, 2015). Although the relationship between these drought tolerance and aphid resistance mechanisms has seldom been explored (Kansman *et al.*, 2022), from a crop breeding perspective it is important to understand the potential for traits to confer cross-tolerance between these two stressors. The mechanisms underpinning effects of water availability on aphid resistance proposed by Leybourne *et al.* (2021) could be examined further for their potential to confer drought tolerance; in Box 1, we illustrate how drought tolerance and aphid resistance traits might interact, and the plant signalling pathways that could communicate cross-tolerance, highlighting potential breeding targets for cross-tolerance.

Conclusions

We highlight that studying the ability of plants to resist aphids under conditions of water restrictions requires consideration that the outcome might be affected by plant genotypic variation in tolerance of drought. Plants may evolve (or be selected through breeding) to express greater drought tolerance, and these traits might also respond to water availability within a generation. The traits and mechanisms underlying aphid resistance and drought tolerance functions may or may not be related but could be subject to trade-offs; understanding their genetic and environmental control is crucial for breeding crops for future climates. Importantly, plant traits that confer aphid tolerance (i.e. compensatory response by plants to damage inflicted by aphids) should be explored for any potential role in plant drought tolerance. As with resistance, both drought and aphid tolerance may have a common molecular and physiological basis and generate cross-tolerance. These views should guide future research in this area.

Acknowledgements

DJL received support from the Alexander von Humboldt Foundation through a Postdoctoral Research Fellowship and from The Royal Commission for the Exhibition of 1851 through an 1851 Research Fellowship. AJK is supported by funding from the Rural and Environment Science and Analytical Services Division of the Scottish Government. CRR is supported by Fondecyt Continuity Fund for Senior Researchers #100462 from University of Talca.

Author contributions

CCR, PEG, AJK, and DJL: conceptualization and writing the manuscript; CCR and DJL: designing the figures.

Conflict of interest

The authors declare no competing interests.

Funding

The Alexander von Humboldt Foundation: Postdoctoral Research Fellowship 'ALAN', the Royal Commission for the Exhibition of 1851 (RF-2022-100004), Rural and Environment Science and Analytical Services Division of the Scottish Government (JHI-A1-2 Integrated Crop Protection), and the National Fund for Scientific and Technological Development, FONDECYT-Chile (grant 1210908 to PEG).

Keywords: Aphid, crop, drought, resistance, susceptible, tolerance, trade-off.

References

- Agrawal AA, Fishbein M. 2008. Phylogenetic escalation and decline of plant defense strategies. *Proceedings of the National Academy of Sciences, USA* **105**, 10057–10060.
- Batyrshina ZS, Shavit R, Yaakov B, Bocobza S, Tzin V. 2022. The transcription factor TaMYB31 regulates the benzoxazinoid biosynthetic pathway in wheat. *Journal of Experimental Botany* **73**, 5634–5649.
- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F. 2012. Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365–377.
- Benkeblia N. 2022. Insights on fructans and resistance of plants to drought stress. *Frontiers in Sustainable Food Systems* **6**, 61.
- Buckley TN. 2019. How do stomata respond to water status? *New Phytologist* **224**, 21–36.
- Denison R. 2012. *Darwinian agriculture: how understanding evolution can improve agriculture*. Princeton, NJ: Princeton University Press.
- Dixon AFG. 1998. *Aphid ecology: an optimization approach*. London: Chapman & Hall.
- Escudero-Martinez CM, Morris JA, Hedley PE, Bos JIB. 2017. Barley transcriptome analyses upon interaction with different aphid species identify thionins contributing to resistance. *Plant, Cell & Environment* **40**, 2628–2643.
- Fang Y, Xiong L. 2015. General mechanisms of drought response and their application in drought resistance improvement in plants. *Cellular and Molecular Life Sciences* **72**, 673–689.
- Foyer CH, Rasool B, Davey JW, Hancock RD. 2016. Cross-tolerance to biotic and abiotic stresses in plants: a focus on resistance to aphid infestation. *Journal of Experimental Botany* **67**, 2025–2037.
- Herms DA, Mattson WJ. 1992. The dilemma of plants: to grow or defend. *The Quarterly Review of Biology* **67**, 283–335.
- Kansman JT, Basu S, Casteel CL, Crowder DW, Lee BW, Nihraz CT, Finke DL. 2022. Plant water stress reduces aphid performance: exploring mechanisms driven by water stress intensity. *Frontiers in Ecology and Evolution* **10**, 288.
- Leimu R, Koricheva J. 2006. A meta-analysis of tradeoffs between plant tolerance and resistance to herbivores: combining the evidence from ecological and agricultural studies. *Oikos* **112**, 1–9.

- Leybourne DJ, Preedy KF, Valentine TA, Bos JIB, Karley AJ.** 2021. Drought has negative consequences on aphid fitness and plant vigor: insights from a meta-analysis. *Ecology and Evolution* **11**, 11915–11929.
- Leybourne DJ, Valentine TA, Binnie K, Taylor A, Karley AJ, Bos JIB.** 2022. Drought stress increases the expression of barley defence genes with negative consequences for infesting cereal aphids. *Journal of Experimental Botany* **73**, 2238–2250.
- Leybourne DJ, Valentine TA, Robertson JAH, Pérez-Fernández E, Main AM, Karley AJ, Bos JIB.** 2019. Defence gene expression and phloem quality contribute to mesophyll and phloem resistance to aphids in wild barley. *Journal of Experimental Botany* **70**, 4011–4026.
- Luo R, Gilbert B.** 2022. Timing of short-term drought structures plant–herbivore dynamics. *Oikos* **2022**, e08860.
- Mitchell C, Brennan RM, Graham J, Karley AJ.** 2016. Plant defense against herbivorous pests: exploiting resistance and tolerance traits for sustainable crop protection. *Frontiers in Plant Science* **7**, 1132.
- Niemeyer HM.** 2009. Hydroxamic acids derived from 2-hydroxy-2H-1,4-benzoxazin-3(4H)-one: key defense chemicals of cereals. *Journal of Agricultural and Food Chemistry* **57**, 1677–1696.
- Osakabe Y, Osakabe K, Shinozaki K, Tran L-SP.** 2014. Response of plants to water stress. *Frontiers in Plant Science* **5**, 86.
- Pearse IS, Aguilar J, Schroder J, Strauss SY.** 2017. Macroevolutionary constraints to tolerance: trade-offs with drought tolerance and phenology, but not resistance. *Ecology* **98**, 2758–2772.
- Peterson RKD, Varella AC, Higley LG.** 2017. Tolerance: the forgotten child of plant resistance. *PeerJ* **5**, e3934.
- Quandahor P, Lin C, Gou Y, A. Coulter J, Liu C.** 2019. Leaf morphological and biochemical responses of three potato (*Solanum tuberosum* L.) cultivars to drought stress and aphid (*Myzus persicae* Sulzer) infestation. *Insects* **10**, 435.
- Sadras VO, Fereres E, Borrás L, Moreno AGE, Araus JL, Fereres A.** 2020. Aphid resistance: an overlooked ecological dimension of nonstructural carbohydrates in cereals. *Frontiers in Plant Science* **11**, 937.
- Sadras V, Vazquez C, Garzo E, Moreno A, Medina S, Taylor J, Fereres A.** 2021. The role of plant labile carbohydrates and nitrogen on wheat–aphid relations. *Scientific Reports* **11**, 12529.
- Saska P, Skuhrovec J, Platková H, Kosová K, Tylová E, Tuan S-J, Vítámvás P.** 2022. Response of the spring wheat–cereal aphid system to drought: support for the plant vigour hypothesis. *Journal of Pest Science*. doi:10.1007/s10340-022-01514-3.
- Saska P, Skuhrovec J, Tylová E, Platková H, Tuan S-J, Hsu Y-T, Vítámvás P.** 2021. Leaf structural traits rather than drought resistance determine aphid performance on spring wheat. *Journal of Pest Science* **94**, 423–434.
- Schneider L, Rebetez M, Rasmann S.** 2022. The effect of climate change on invasive crop pests across biomes. *Current Opinion in Insect Science* **50**, 100895.
- Strauss SY, Agrawal AA.** 1999. The ecology and evolution of plant tolerance to herbivory. *Trends in Ecology & Evolution* **14**, 179–185.
- Tardieu F.** 2022. Different avenues for progress apply to drought tolerance, water use efficiency and yield in dry areas. *Current Opinion in Biotechnology* **73**, 128–134.
- Valim JOS, Teixeira NC, Santos NA, Oliveira MGA, Campos WG.** 2016. Drought-induced acclimatization of a fast-growing plant decreases insect performance in leaf-chewing and sap-sucking guilds. *Arthropod-Plant Interactions* **10**, 351–363.
- Zhou S, Richter A, Jander G.** 2018. Beyond defense: multiple functions of benzoxazinoids in maize metabolism. *Plant and Cell Physiology* **59**, 1528–1537.